

with tobacco amblyopia and nutritional (tropical) amblyopia at risk in that hydroxocobalamin, and not cyanocobalamin, is a powerful cyanide antagonist. If such patients are given cyanocobalamin, the diagnosis may be questioned and the patient be thus condemned to a life of poor sight.

Patients with tobacco amblyopia who have normal serum vitamin B12 levels need not continue therapy with systemic hydroxocobalamin once their visual acuity and visual fields have returned to normal providing they abstain from further smoking. However, patients who have low serum vitamin B12 levels and evidence of defective vitamin B12 absorption will need to continue indefinitely with hydroxocobalamin as will all patients with proven pernicious anaemia who are at risk of developing optic neuropathy if they are, or later become, smokers.

I would remind Dr Lederle that besides being present in tobacco-smoke and alcohol, cyanide has a worldwide distribution in the plant kingdom as well as being present in industrial waste. There may thus come a time when more widespread chronic cyanide neurotoxicity occurs from a dietary source in persons with a genetic or acquired error of cyanide or vitamin B12 metabolism and that such patients will only respond to treatment with systemic hydroxocobalamin⁵.

A G FREEMAN

Meadow Rise
3 Lakeside
Swindon SN3 1QE

References

- 1 Matthews DM. Distribution of cobalamins in the animal body. In: Zagalak B, Friedrich W, eds. *Vitamin B12: Proceedings of the 3rd European Symposium*. Berlin: De Gruyter, 1979:681-94
- 2 *Drugs and therap Bull* 1984;22:43-4
- 3 Chipping PM. Vitamin B12 deficiency. *Prescribers' J* 1988;28:117-24
- 4 *British National Formulary*. March 1991:296-7
- 5 Freeman AG. Chronic cyanide intoxication. *BMJ* 1981; 282:1321

I should like to correct an error in the paper by Dr Freeman (November 1992 *JRSM*, pp 686-7). Since at least the fifth *World Health Organization Model List of Essential Drugs*, published in 1988, hydroxocobalamin, not cyanocobalamin, has been the drug included.

MARCUS M REIDENBERG

The New York Hospital
Cornell Medical Center
525 East 68th Street,
Box 70, New York, NY 10021, USA

The author replies below:

I do not dispute the data supplied by Professor Reidenberg but this, with respect, was not the point at issue.

In 1970 we presented a case for withdrawal of cyanocobalamin in favour of hydroxocobalamin for therapeutic use. As no action was taken by the manufacturers, we again asked in 1978 'why has cyanocobalamin not been withdrawn?'. This view received widespread support. Yet, until 1988, the World Health Organization's Committee on the selection of essential drugs continued to list cyanocobalamin to the exclusion of hydroxocobalamin, thus placing an incalculable number of patients

with tobacco and nutritional amblyopia and optic neuropathy in pernicious anaemia at risk.

A G FREEMAN

Meadow Rise
3 Lakeside
Swindon N3 1QE

He/she

I refer to the correspondence in the *Journal* (by Ellis and Ridgway (April 1992 *JRSM*, p 246), Ellerby and Lindsey (September 1992 *JRSM*, p 586) and Jackson (October 1992 *JRSM*, p 651) regarding the necessity for a sex-neutral personal pronoun in English. The solutions proposed by Ellerby (s/he) and Ellis (hshe) could be used in writing, but their spoken pronunciation is not obvious. Lindsey's suggestions are syntactically ingenious, but unpronounceable, and at odds with the historical tendency of English usage to eliminate case differences.

I therefore propose the neologism 'esh' as the case-insensitive, sex-neutral personal pronoun so clearly needed in late twentieth century English, and look forward to its enthusiastic and trend-setting adoption by *JRSM* in place of the inflammatory sex-specific pronouns and of the awkward circumlocutions often used to avoid them.

SIDNEY N KAHN

1425 Grasshopper Road
Huntingdon Valley, PA 19006, USA

Nitrous oxide at homeopathic doses

I would like to comment on some of the issues raised by Dr R Schoental in her recent letter (September 1992 *JRSM*, p 587). A number of studies on the cognitive consequences of trace concentrations of nitrous oxide (in the region of 500 ppm), have revealed an absence of effects, although mood was not specifically examined¹. None the less, analgesic concentrations of nitrous oxide can be antidepressant in certain subjects². The effects of the homeopathic concentrations alluded to by Dr Schoental, have, as far as I am aware not been examined for their mood altering effects. However, there is no evidence that concentrations of nitrous oxide as low as in the ppb range can have any effect on folate and cobalamin metabolism^{3,4}.

There is another point; nitrous oxide as an anaesthetic is not especially used for paediatric dentistry. Today, as previously, it is employed, as an anaesthetic for all forms of surgery, both general and dental. None the less, it is also used in dentistry, obstetrics and for treating addictive withdrawal states at analgesic doses in which there is no loss of consciousness^{2,5}.

M A GILLMAN

South African Brain
Research Institute
6 Campbell Street
Waveley 2090 Johannesburg
South Africa

References

- 1 Frost EAM. In: E I Eger II (ed). *Central nervous effects of nitrous oxide in nitrous oxide/N₂O*. New York: Elsevier, 1985:157-76
- 2 Gillman MA, Lichtigfeld FJ. Analgesic nitrous oxide in neuropsychiatry: past present and future. *Int J Neurosci* 1989;49:75-81
- 3 Nunn JF, Chananin I. In: E I Eger II (ed). *Nitrous oxide inactivates methionine synthetase in nitrous oxide/N₂O*. New York: Elsevier, 1985:211-33

- 4 Gillman MA. Haematological changes caused by nitrous oxide, cause for concern? Editorial. *Br J Anaesth* 1987;59:143-5
- 5 Gillman MA, Lichtigfeld FJ. Analgesic nitrous oxide for alcohol withdrawal: a critical appraisal after 10 years' use. *Postgrad Med J* 1990;66:543-6

The author replies below:

I am delighted to learn that Dr Gillman is interested in the effects of pharmacological concentrations of nitrous oxide in neuropsychiatry. He may be able to establish the relation of the concentrations of this compound in the atmosphere to the factors that determine human mood, a problem which I am not able to undertake, but appears to warrant appropriate studies.

It may be worth noting that in a study of the global and seasonal variability of the composition of stratosphere and of the mesosphere by the experimental meteorological satellite Nimbus 7 (launched in 1978), only the data for nitrous oxide (N₂O) appeared to differ from those expected from a prediction model¹.

R SCHOENTAL

15 Birdhurst Court,
Woodcote Road,
Wallington,
Surrey SM6 0PG

References

- 1 Barnett JJ, Corney M, Murphy AK, Jones RL, Rodgers CD, Taylor FW, *et al*. Global and seasonal variability of the temperature and composition of the middle atmosphere. *Nature* 1985; 313:439-43

Post anoxic action myoclonus

We read with interest the recent report of Rustom and Fiskén (December 1992 *JRSM*, p 761) on post anoxic action myoclonus. Increased awareness of this condition, on both the intensive care unit¹ and general medical wards, is to be welcomed as a prerequisite for appropriate treatment, but we would take issue with some of the authors' statements regarding its pathophysiology and treatment.

Despite low levels of CSF metabolites of 5HT, not all patients with this syndrome respond to serotonergic drugs². Furthermore, although, as stated, clonazepam is the drug of choice, the evidence available is against its action being at 5HT receptors³, indeed its effects (reduced 5HT utilization in brain) are exactly the opposite of those predicted from the beneficial effects of 5-hydroxytryptophan in human myoclonic disorders³. Hence, an action at GABA receptor complexes seems more likely³, although why clonazepam should be more efficacious than other benzodiazepines is unclear. CSF GABA levels are low in post anoxic myoclonus, and GABA agonists such as baclofen have occasionally been reported of benefit⁴.

Contrary to the implications of Rustom and Fiskén, clonazepam does not always produce a dramatic therapeutic response; other agents (a wide variety of which have been tried⁵) may need to be added, especially sodium valproate². Another drug which may be useful is piracetam (2-oxo-1-pyrrolidine acetamide): it is claimed to be of value in some patients with cognitive deficits, especially of hypoxic origin, and on the basis of its hypothesized effects on energy metabolism has been tried in post anoxic myoclonus. Its chemical structure is similar to GABA but its precise mechanism of action is unknown⁶. We recently treated a 56-year-old man who suffered anoxic brain damage following a respiratory arrest during a severe asthma attack; his myoclonic movements were not well controlled by clonazepam (10 mg/day) and sodium valproate (4 mg/day), but addition of piracetam (UCB SA Pharmaceuticals, Brussels, Belgium; 7.2 g/day) produced considerable amelioration with marked improvement in the performance of the activities of daily living. Other reports of the efficacy of piracetam have appeared^{2,7-9}, and it seems of particular use in patients with evidence of cortical (as opposed to reticular) reflex myoclonus.

A combination of drugs may therefore be necessary in severe post anoxic myoclonus². Since the site, severity, and extension of hypoxia-induced damage is variable between patients, differing responsiveness to drugs is not surprising and suggests the need for an empirical approach to therapy.

ANDREW J LARNER

M TOM E HEAFIELD

University Department
of Clinical Neurology
The University of Birmingham,
Edgbaston, Birmingham B15 2TH

References

- 1 Harper SJ, Wilkes RG. Posthypoxic myoclonus (the Lance-Adams syndrome) in the intensive care unit. *Anaesthesia* 1991;46:199-201
- 2 Obeso JA, Artieda J, Rothwell JC, Day B, Thompson P, Marsden CD. The treatment of severe action myoclonus. *Brain* 1989;112:765-77
- 3 Jenner P, Pratt JA, Marsden CD. Mechanism of action of clonazepam in myoclonus in relation to effects of GABA and 5-HT. *Adv Neurol* 1986;43:629-43
- 4 Coletti A, Mandelli A, Minoli G, Tredici G. Post-anoxic action myoclonus (Lance-Adams syndrome) treated with levodopa and GABAergic drugs. *J Neurol* 1980;223:67-70
- 5 Fahn S. Posthypoxic action myoclonus: literature review update. *Adv Neurol* 1985;43:157-69
- 6 Tacconi MT, Wurtman RJ. Piracetam: physiological disposition and mechanism of action. *Adv Neurol* 1986;43:675-85
- 7 Terwinghe G, Daumerie J, Nicaise C, Rosillon O. Effet thérapeutique du piracetam dans un cas de myoclonie d'action postanoxique. *Acta Neurol Belg* 1978;78:30-6
- 8 Cremieux C, Serratrice G. Myoclonie d'intention et d'action post-anoxique: amélioration par le piracetam. *Nouv Presse Med* 1979;41:3357-8
- 9 Fahn S. Newer drugs for posthypoxic action myoclonus: observations from a well-studied case. *Adv Neurol* 1986;43:197-9